

The small index property of the Fraïssé limit of finite Heyting algebras

Kentarô Yamamoto

Institute of Computer Science, the Czech Academy of Sciences

June 13, 2022

Motivation: Problem of reconstruction

Problem

For a reasonable class \mathcal{K} of (countable) structures and $M, N \in \mathcal{K}$:

$$\text{Aut}(M) \cong \text{Aut}(N) \stackrel{?}{\iff} M \cong N$$

This is a way to precisify the broader question:

Problem

Is it possible to describe a structure *independently* of signatures?

Example: Countable atomless Boolean algebra

Let \mathbb{B} be the countable atomless Boolean algebra.

Fact

For $a, a', b, b' \in \mathbb{B} \setminus \{0, 1\}$, if $a \leq b$ and $a' \leq b'$, then there is $\sigma \in \text{Aut}(\mathbb{B})$ s.t. $\sigma(ab) = a'b'$.

Let B be the domain of \mathbb{B} , and its automorphism group $G \subseteq {}^B B$ is given as a **permutation group**.

- 1 There are exactly two fixed points $\{0, 1\}$ of the action.
- 2 Consider the action $G \curvearrowright B^2$.
- 3 There are three orbits of the action: $\{(0, 1)\}, R, S$.
- 4 Exactly one of R, S (say, R) is a reflexive binary relation.
- 5 Let $\leq = R \cup (\{0\} \times B) \cup (B \times \{1\})$.

Ultrahomogeneity

Observation

It helps reconstruction to have many automorphisms.

Definition

A countable structure M is **ultrahomogeneous** if every isomorphism between finitely generated substructures of M extends to an automorphism of M .

Example

The countable atomless Boolean algebra is ultrahomogeneous.

Different variants of reconstruction

There are a few ways to interpret the LHS of

$$\text{Aut}(M) \cong \text{Aut}(N) \stackrel{?}{\iff} M \cong N \quad (1)$$

① Isomorphic as permutation groups

② Isomorphic as **topological** groups

Here $\text{Aut}(M)$ has the **pointwise convergence topology**, i.e., the induced topology as a subset of ${}^\omega\omega$.

③ Isomorphic as abstract groups

Automorphism groups lose information as you go further down in the list.

Fact

If M and N are ω -categorical, then (1) holds as far as the isomorphism as topological groups is concerned.

Subgroups of small indices

Fact

For a countable structure M and subgroup $H \leq \text{Aut}(M)$, the following are equivalent:

- 1 H is open.
- 2 $(G : H) \leq \aleph_0$.
- 3 $(G : H) < 2^{\aleph_0}$.

Definition

A topological group has the **small index property**

\iff every subgroup of index less than 2^{\aleph_0} is open.

A countable structure M has the s.i.p. $\iff \text{Aut}(M)$ does.

If $\text{Aut}(M)$ has the small index property, its topology can be recovered from its abstract group structure only.

Which structures have the small index property?

Empirically, countable ultrahomogeneous structures fall into exactly one of the following classes:

- 1 Those known to have the small index property (e.g., the countable atomless Boolean algebra)
- 2 Structures made up so they wouldn't have the small index property
- 3 Structures not known to have the small index property (e.g., the universal ultrahomogeneous poset)

Fraïssé limit of finite Heyting algebras

The existence of the following follows from a general fact model theory and the amalgamation property of Heyting algebras (e.g., Maksimova 1977):

Definition

The Fraïssé limit L of finite Heyting algebras is the unique countable ultrahomogeneous structure into which every finite nontrivial Heyting algebra embeds.

Informally it lies halfway between the universal ultrahomogeneous poset and the countable atomless Boolean algebra.

Theorem

The Fraïssé limit L of finite Heyting algebras has the small index property.

Definition

A Polish group G has **ample generic** elements

\iff for each $n < \omega$ the action of G on G^n defined by

$$g \cdot (\dots, h, \dots) = (\dots, ghg^{-1}, \dots)$$

has a comeager orbit.

Theorem

If $\text{Aut}(M)$ has ample generics, then it has the strong index property (Kechris and Rosendal 2006)

It appears that if M has a “nontrivial” order, this method cannot be used.

Our method I

Instead, we adapted the old argument by Truss 1989 establishing the small index property of the countable Boolean algebra.

Theorem

The class of nontrivial finite Heyting algebras has the *superamalgamation property*: Given finite Heyting algebras A_0, A_1, A_2 with $A_0 \subseteq (A_1 \cup A_2)$, there exists a finite Heyting algebra A containing $A_1 \cup A_2$ such that for $(i, j) = (1, 2), (2, 1)$

$$\forall (a_i \in A_i) \forall (a_j \in A_j) [a_i \leq a_j \implies \exists a_0 \in A_0 \ a_i \leq a_0 \leq a_j]$$

(Maksimova 1977)





Theorem

$\text{Aut}(M)$ is simple. (Yamamoto 202?)

Our argument also uses the following fact:

Lemma

Let X be the Esakia space of L , and consider the induced action of G on X . Let $Y \subseteq X$ be nonempty and finite. Then the G -orbit $G \cdot Y$ of Y in the natural action $G \curvearrowright \mathcal{P}(X)$ is of cardinality continuum.

-  Kechris, A. and C. Rosendal (2006). “Turbulence, Amalgamation, and Generic Automorphisms of Homogeneous Structures”. In: *Proceedings of the London Mathematical Society*.
-  Maksimova, L. L. (Nov. 1977). “Craig’s Theorem in Superintuitionistic Logics and Amalgamable Varieties of pseudo-Boolean Algebras”. In: *Algebra and Logic* 16.6, pp. 427–455.
-  Truss, J. K. (1989). “Infinite Permutation Groups II. Subgroups of Small Index”. In: *Journal of Algebra* 120.2, pp. 494 –515. ISSN: 0021-8693.
-  Yamamoto, K. (202?). “The Automorphism group of the Fraïssé Limit of Heyting Algebras”. In: *The Journal of Symbolic Logic*.